

# Agenda

## Chebyshev Approximation (I)

- Galerkin Approach

- Collocation Approach

- Reciprocal Relations & Economization of Power Series

## Chebyshev Approximation (II)

- Spectral Differentiation

- Differentiation in Real Space

## Implementation of Boundary Conditions

- Galerkin Approach & Basis Recombination

- Galerkin Approach & Tau Method

- Collocation Method

- ▶ Consider an approximation of  $u \in L^2_\omega(I)$  in terms of a **TRUNCATED CHEBYSHEV SERIES**  $u_N(x) = \sum_{k=0}^N \hat{u}_k T_k(x)$
- ▶ Cancel the projections of the residual  $R_N = u - u_N$  on the  $N + 1$  first basis function (i.e., the Chebyshev polynomials)

$$(R_N, T_l)_\omega = \int_{-1}^1 \left( u T_l \omega - \sum_{k=0}^N \hat{u}_k T_k T_l \omega \right) dx = 0, \quad l = 0, \dots, N$$

- ▶ Taking into account the orthogonality condition, expressions for the Chebyshev expansions coefficients are obtained

$$\hat{u}_k = \frac{2}{\pi c_k} \int_{-1}^1 u T_k \omega dx,$$

which can be evaluated using, e.g., the

**GAUSS-LOBATTO-CHEBYSHEV QUADRATURES** .

- ▶ **QUESTION** — What happens on the boundary?

## Theorem

Let  $P_N : L^2_\omega(I) \rightarrow \mathbb{P}_N$  be the orthogonal projection on the subspace  $\mathbb{P}_N$  of polynomials of degree  $\leq N$ . For all  $\mu$  and  $\sigma$  such that  $0 \leq \mu \leq \sigma$ , there exists a constant  $C$  such that

$$\|u - P_N u\|_{\mu, \omega} < CN^{e(\mu, \sigma)} \|u\|_{\sigma, \omega}$$

$$\text{where } e(\mu, \sigma) = \begin{cases} 2\mu - \sigma - \frac{1}{2} & \text{for } \mu > 1, \\ \frac{3}{2}\mu - \sigma & \text{for } 0 \leq \mu \leq 1 \end{cases}$$

“Philosophy” of the proof.

1. First establish continuity of the mapping  $u \rightarrow \tilde{u}$ , where  $\tilde{u}(\theta) = u(\cos(\theta))$ , from the weighted Sobolev space  $H^m_\omega(I)$  into the corresponding periodic Sobolev space  $H^m_p(-\pi, \pi)$
2. Then leverage analogous approximation error bounds established for the case of trigonometric basis functions

- ▶ Consider an approximation of  $u \in L^2_\omega(I)$  in terms of a truncated Chebyshev series (expansion coefficients as the unknowns)

$$u_N(x) = \sum_{k=0}^N \hat{u}_k T_k(x)$$

- ▶ Cancel the residual  $R_N = u - u_N$  on the set of GAUSS-LOBATTO-Chebyshev collocation points  $x_j$ ,  $j = 0, \dots, N$  (one could choose other sets of collocation points as well)

$$u(x_j) = \sum_{k=0}^N \hat{u}_k T_k(x_j), \quad j = 0, \dots, N$$

- ▶ Noting that  $T_k(x_j) = \cos\left(k \cos^{-1}\left(\cos\left(\frac{j\pi}{N}\right)\right)\right) = \cos\left(k \frac{j\pi}{N}\right)$  and denoting  $u_j \triangleq u(x_j)$  we obtain

$$u_j = \sum_{k=0}^N \hat{u}_k \cos\left(k \frac{\pi j}{N}\right), \quad j = 0, \dots, N$$

- ▶ The above system of equations can be written as  $U = \mathcal{T}\hat{U}$ , where  $U$  and  $\hat{U}$  are vectors of grid values and expansion coefficients, respectively.

- ▶ In fact, the matrix  $\mathcal{T}$  is invertible and

$$[\mathcal{T}^{-1}]_{jk} = \frac{2}{\bar{c}_j \bar{c}_k N} \cos\left(\frac{k\pi j}{N}\right), \quad j, k = 0, \dots, N$$

- ▶ Consequently, the expansion coefficients can be expressed as follows

$$\hat{u}_k = \frac{2}{\bar{c}_k N} \sum_{j=0}^N \frac{1}{\bar{c}_j} u_j \cos\left(\frac{k\pi j}{N}\right) = \frac{2}{\bar{c}_k N} \sum_{j=0}^N \frac{1}{\bar{c}_j} u_j \Re\left[e^{i\left(\frac{k\pi j}{N}\right)}\right], \quad k = 0, \dots, N$$

Note that this expression is nothing else than the **COSINE TRANSFORM** of  $U$  which can be very efficiently evaluated using a **COSINE FFT**

- ▶ The same expression can be obtained by
  - ▶ multiplying each side of  $u_j = \sum_{k=0}^N \hat{u}_k T_k(x_j)$  by  $\frac{T_l(x_j)}{\bar{c}_j}$
  - ▶ summing the resulting expression from  $j = 0$  to  $j = N$
  - ▶ using the **DISCRETE ORTHOGONALITY RELATION**

$$\frac{\pi}{N} \sum_{j=0}^N \frac{1}{\bar{c}_j} T_k(\tilde{\xi}_j) T_l(\tilde{\xi}_j) = \frac{\pi \bar{c}_k}{2} \delta_{kl}$$

- ▶ Note that the expression for the **DISCRETE CHEBYSHEV TRANSFORM**

$$\hat{u}_k = \frac{2}{c_k N} \sum_{j=0}^N \frac{1}{c_j} u_j \cos\left(\frac{k\pi j}{N}\right), \quad k = 0, \dots, N$$

can also be obtained by using the **Gauss-Lobatto-Chebyshev** quadrature to approximate the continuous expressions

$$\hat{u}_k = \frac{2}{\pi c_k} \int_{-1}^1 u T_k \omega dx, \quad k = 0, \dots, N,$$

Such an approximation is **EXACT** for  $u \in \mathbb{P}_N$

- ▶ Analogous expressions for the Discrete Chebyshev Transforms can be derived for other set of collocation points (Gauss, Gauss-Radau)
- ▶ Note similarities with respect to the case periodic functions and the Discrete Fourier Transform

- ▶ As was the case with Fourier spectral methods, there is a very close connection between **COLLOCATION-BASED INTERPOLATION** and **GALERKIN APPROXIMATION**
- ▶ **DISCRETE CHEBYSHEV TRANSFORM** can be associated with an **INTERPOLATION OPERATOR**  $P_C : C^0(I) \rightarrow \mathbb{R}^N$  defined such that  $(P_C u)(x_j) = u(x_j)$ ,  $j = 0, \dots, N$  (where  $x_j$  are the Gauss-Lobatto collocation points)

### Theorem

Let  $s > \frac{1}{2}$  and  $\sigma$  be given and  $0 \leq \sigma \leq s$ . There exists a constant  $C$  such that

$$\|u - P_C u\|_{\sigma, \omega} < CN^{2\sigma-s} \|u\|_{s, \omega}$$

for all  $u \in H_{\omega}^s(I)$ .

### Outline of the Proof.

Changing the variables to  $\tilde{u}(\theta) = u(\cos(\theta))$  we convert this problem to a problem already analyzed in the context of the Fourier interpolation for periodic functions □

- Relation between the **GALERKIN** and **COLLOCATION** coefficients, i.e.,

$$\hat{u}_k^e = \frac{2}{\pi c_k} \int_{-1}^1 u(x) T_k(x) \omega(x) dx, \quad k = 0, \dots, N$$

$$\hat{u}_k^c = \frac{2}{\bar{c}_k N} \sum_{j=0}^N \frac{1}{\bar{c}_j} u_j \cos\left(\frac{k\pi j}{N}\right), \quad k = 0, \dots, N$$

- Using the representation  $u(x) = \sum_{l=0}^{\infty} \hat{u}_l^e T_l(x)$  in the latter expression and invoking the discrete orthogonality relation we obtain

$$\begin{aligned} \hat{u}_k^c &= \frac{2}{\bar{c}_k N} \sum_{l=0}^N \hat{u}_l^e \left[ \sum_{j=0}^N \frac{1}{\bar{c}_j} T_k(x_j) T_l(x_j) \right] + \frac{2}{\bar{c}_k N} \sum_{l=N+1}^{\infty} \hat{u}_l^e \left[ \sum_{j=0}^N \frac{1}{\bar{c}_j} T_k(x_j) T_l(x_j) \right], \\ &= \hat{u}_k^e + \frac{2}{\bar{c}_k N} \sum_{l=N+1}^{\infty} \hat{u}_l^e C_{kl} \end{aligned}$$

$$\text{where } C_{kl} = \sum_{j=0}^N \frac{1}{\bar{c}_j} T_k(x_j) T_l(x_j) = \sum_{j=0}^N \frac{1}{\bar{c}_j} \cos\left(\frac{kj\pi}{N}\right) \cos\left(\frac{lj\pi}{N}\right)$$

$$= \frac{1}{2} \sum_{j=0}^N \frac{1}{\bar{c}_j} \left[ \cos\left(\frac{k-l}{N}j\pi\right) + \cos\left(\frac{k+l}{N}j\pi\right) \right]$$



- ▶ Using the identity

$$\sum_{j=0}^N \cos\left(\frac{pj\pi}{N}\right) = \begin{cases} N+1, & \text{if } p = 2mN, \quad m = 0, \pm 1, \pm 2, \dots \\ \frac{1}{2}[1 + (-1)^p] & \text{otherwise} \end{cases}$$

we can calculate  $C_{kl}$  which allows us to express the relation between the Galerkin and collocation coefficients as follows

$$\hat{u}_k^c = \hat{u}_k^e + \frac{1}{\bar{c}_k} \left[ \sum_{\substack{m=1 \\ 2mN > N-k}}^{\infty} \hat{u}_{k+2mN}^e + \sum_{\substack{m=1 \\ 2mN > N+k}}^{\infty} \hat{u}_{-k+2mN}^e \right]$$

- ▶ The terms in square brackets represent the **ALIASING ERRORS**. Their origin is precisely the same as in the Fourier (pseudo)-spectral method.
- ▶ Aliasing errors can be removed using the **3/2 APPROACH** in the same way as in the Fourier (pseudo)-spectral method

- ▶ expressing the first  $N$  Chebyshev polynomials as functions of  $x^k$ ,  
 $k = 1, \dots, N$

$$T_0(x) = 1,$$

$$T_1(x) = x,$$

$$T_2(x) = 2x^2 - 1,$$

$$T_3(x) = 4x^3 - 3x,$$

$$T_4(x) = 8x^4 - 8x^2 + 1$$

which can be written as  $V = \mathbb{K}X$ , where  $[V]_k = T_k(x)$ ,  $[X]_k = x^k$ , and  $\mathbb{K}$  is a **LOWER-TRIANGULAR** matrix

- ▶ Solving this system (trivially!) results in the following **RECIPROCAL RELATIONS**

$$1 = T_0(x),$$

$$x = T_1(x),$$

$$x^2 = \frac{1}{2}[T_0(x) + T_2(x)],$$

$$x^3 = \frac{1}{4}[3T_1(x) + T_3(x)],$$

$$x^4 = \frac{1}{8}[3T_0(x) + 4T_2(x) + T_4(x)]$$

- ▶ Find the best polynomial approximation of order 3 of  $f(x) = e^x$  on  $[-1, 1]$
- ▶ Construct the (Maclaurin) expansion

$$e^x = 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \dots$$

- ▶ Rewrite the expansion in terms of **CHEBYSHEV POLYNOMIALS** using the reciprocal relations

$$e^x = \frac{81}{64} T_0(x) + \frac{9}{8} T_1(x) + \frac{13}{48} T_2(x) + \frac{1}{24} T_3(x) + \frac{1}{192} T_4(x) + \dots$$

- ▶ Truncate this expansion to the 3<sup>rd</sup> order and translate the expansion back to the  $x^k$  representation
- ▶ Truncation error is given by the magnitude of the first truncated term; Note that the **CHEBYSHEV EXPANSION COEFFICIENTS** are much smaller than the corresponding **TAYLOR EXPANSION COEFFICIENTS** !
- ▶ How is it possible – the same number of expansion terms, but higher accuracy?

- ▶ Assume function approximation in the form  $u_N(x) = \sum_{k=0}^N \hat{u}_k T_k(x)$
- ▶ First, note that CHEBYSHEV PROJECTION and DIFFERENTIATION do not commute, i.e.,  $P_N(\frac{du}{dx}) \neq \frac{d}{dx}(P_N u)$
- ▶ Sequentially applying the recurrence relation  $2T_k = \frac{T'_{k+1}}{k+1} - \frac{T'_{k-1}}{k-1}$  we obtain

$$T'_k(x) = 2k \sum_{p=0}^K \frac{1}{c_{k-1-2p}} T_{k-1-2p}(x), \quad \text{where } K = \left\lfloor \frac{k-1}{2} \right\rfloor$$

- ▶ Consider the first derivative

$$u'_N(x) = \sum_{k=0}^N \hat{u}_k T'_k(x) = \sum_{k=0}^N \hat{u}_k^{(1)} T_k(x)$$

where, using the above expression for  $T'_k(x)$ , we obtain the expansion coefficients as

$$\hat{u}_k^{(1)} = \frac{2}{c_k} \sum_{\substack{p=k+1 \\ (p+k) \text{ odd}}}^N p \hat{u}_p, \quad k = 0, \dots, N-1, \quad \text{and} \quad \hat{u}_N^{(1)} = 0$$

- ▶ Spectral differentiation (with the expansion coefficients as unknowns) can thus be written as

$$\hat{U}^{(1)} = \hat{\mathbb{D}}\hat{U},$$

where  $\hat{U} = [\hat{u}_0 \dots, \hat{u}_N]^T$ ,  $\hat{U}^{(1)} = [\hat{u}_0^{(1)} \dots, \hat{u}_N^{(1)}]^T$ , and  $\hat{\mathbb{D}}$  is an **UPPER-TRIANGULAR** matrix with entries deduced based on the previous expression

- ▶ For the second derivative one obtains similarly

$$u_N''(x) = \sum_{k=0}^N \hat{u}_k^{(2)} T_k(x)$$

$$\hat{u}_k^{(2)} = \frac{1}{c_k} \sum_{\substack{p=k+2 \\ (p+k) \text{ even}}}^N p(p^2 - k^2) \hat{u}_p, \quad k = 0, \dots, N-2$$

and  $\hat{u}_N^{(2)} = \hat{u}_{N-1}^{(2)} = 0$

- ▶ **QUESTION** — What is the structure of the second-order differentiation matrix?

- Assume the function  $u(x)$  is approximated in terms of its nodal values, i.e.,

$$u(x) \cong u_N(x) = \sum_{j=0}^N u(x_j) C_j(x),$$

where  $\{x_j\}$  are the **GAUSS-LOBATTO-CHEBYSHEV** points and  $C_j(x)$  are the associated **CARDINAL FUNCTIONS**

$$C_j(x) = (-1)^{j+1} \frac{(1-x^2)}{c_j N^2 (x-x_j)} \frac{dT_N(x)}{dx} = \frac{2}{N p_j} \sum_{m=0}^N \frac{1}{p_m} T_m(x_j) T_m(x),$$

where

$$p_j = \begin{cases} 2 & \text{for } j = 0, N, \\ 1 & \text{for } j = 1, \dots, N-1, \end{cases} \quad c_j = \begin{cases} 2 & \text{for } j = N, \\ 1 & \text{for } j = 0, \dots, N-1 \end{cases}$$

- The **DIFFERENTIATION MATRIX**  $\mathbb{D}^{(p)}$  relating the nodal values of the  $p$ -th derivative  $u_N^{(p)}$  to the nodal values of  $u$  is obtained by differentiating the cardinal function appropriate number of times

$$u_N^{(p)}(x_j) = \sum_{k=0}^N \frac{d^{(p)} C_k(x_j)}{dx^{(p)}} u(x_k) = \sum_{k=0}^N d_{jk}^{(p)} u(x_k), \quad j = 0, \dots, N$$

- ▶ Expressions for the entries of the **DIFFERENTIATION MATRIX**  $d_{jk}^{(1)}$  at the the **GAUSS-LOBATTO-CHEBYSHEV** collocation points

$$d_{jk}^{(1)} = \frac{\bar{c}_j}{\bar{c}_k} \frac{(-1)^{j+k}}{x_j - x_k}, \quad 0 \leq j, k \leq N, j \neq k,$$

$$d_{jj}^{(1)} = -\frac{x_j}{2(1-x_j^2)}, \quad 1 \leq j \leq N-1,$$

$$d_{00}^{(1)} = -d_{NN}^{(1)} = \frac{2N^2 + 1}{6},$$

- ▶ Thus in the matrix (operator) notation

$$U^{(1)} = \mathbb{D}U$$

- ▶ Note that **ROWS** of the differentiation matrix  $\mathbb{D}$  are in fact equivalent to  $N$ -th order asymmetric finite-difference formulas on a nonuniform grid; in other words, spectral differentiation using nodal values as unknowns is equivalent to finite differences employing **ALL  $N$  GRID POINTS AVAILABLE**

- Expressions for the entries of **SECOND-ORDER DIFFERENTIATION MATRIX**  $d_{jk}^{(2)}$  at the the **GAUSS-LOBATTO-CHEBYSHEV** collocation points ( $U^{(2)} = \mathbb{D}^{(2)} U$ )

$$d_{jk}^{(2)} = \frac{(-1)^{j+k}}{\bar{c}_k} \frac{x_j^2 + x_j x_k - 2}{(1 - x_j^2)(x_j - x_k)^2}, \quad 1 \leq j \leq N-1, 0 \leq k \leq N, j \neq k$$

$$d_{jj}^{(2)} = -\frac{(N^2 - 1)(1 - x_j^2) + 3}{3(1 - x_j^2)^2}, \quad 1 \leq j \leq N-1,$$

$$d_{0k}^{(2)} = \frac{2}{3} \frac{(-1)^k}{\bar{c}_k} \frac{(2N^2 + 1)(1 - x_k) - 6}{(1 - x_k)^2}, \quad 1 \leq k \leq N$$

$$d_{Nk}^{(2)} = \frac{2}{3} \frac{(-1)^{N+k}}{\bar{c}_k} \frac{(2N^2 + 1)(1 + x_k) - 6}{(1 + x_k)^2}, \quad 0 \leq k \leq N-1$$

$$d_{00}^{(2)} = d_{NN}^{(2)} = \frac{N^4 - 1}{15},$$

- Note that 
$$d_{jk}^{(2)} = \sum_{p=0}^N d_{jp}^{(1)} d_{pk}^{(1)}$$

- Interestingly,  $\mathbb{D}^2$  is not a **SYMMETRIC MATRIX** ...



- Consider an **ELLIPTIC BOUNDARY VALUE PROBLEM (BVP)** :

$$\begin{aligned} -\nu u'' + au' + bu &= f, & \text{in } [-1, 1] \\ \alpha_- u + \beta_- u' &= g_- & x = -1 \\ \alpha_+ u + \beta_+ u' &= g_+ & x = 1 \end{aligned}$$

- Chebyshev polynomials do not satisfy homogeneous boundary conditions, hence standard Galerkin approach is not directly applicable.
- **BASIS RECOMBINATION** :

- Convert the BVP to the corresponding form with **HOMOGENEOUS BOUNDARY CONDITIONS**
- Take linear combinations of Chebyshev polynomials to construct a new basis satisfying **HOMOGENEOUS DIRICHLET BOUNDARY CONDITIONS**  
 $\varphi_k(\pm 1) = 0$

$$\varphi_k(x) = \begin{cases} T_k(x) - T_0(x) = T_k - 1, & k - \text{even} \\ T_k(x) - T_1(x), & k - \text{odd} \end{cases}$$

Note that the new basis preserves orthogonality

- ▶ **THE TAU METHOD** (Lanczos, 1938) consists in using a Galerkin approach in which explicit enforcement of the boundary conditions replaces projections on some of the test functions
- ▶ Consider the residual

$$R_N(x) = -\nu u_N'' + au_N' + bu_N - f,$$

where  $u_N(x) = \sum_{k=0}^N \hat{u}_k T_k(x)$

- ▶ Cancel projections of the residual on the first  $N - 2$  basis functions

$$(R_N, T_l)_\omega = \sum_{k=0}^N \left( -\nu \hat{u}_k^{(2)} + a \hat{u}_k^{(1)} + b \hat{u}_k \right) \int_{-1}^1 T_k T_l \omega dx - \int_{-1}^1 f T_l \omega dx, \quad l = 0, \dots, N - 2$$

- ▶ Thus, using orthogonality, we obtain

$$-\nu \hat{u}_k^{(2)} + a \hat{u}_k^{(1)} + b \hat{u}_k = \hat{f}_k, \quad k = 0, \dots, N - 2$$

where  $\hat{f}_k = \int_{-1}^1 f T_k \omega dx$

- ▶ Noting that  $T_k(\pm 1) = (\pm 1)^k$  and  $T'_k(\pm 1) = (\pm 1)^{k+1}k^2$ , the **BOUNDARY CONDITIONS** are enforced by supplementing the residual equations with

$$\sum_{k=0}^N (-1)^k (\alpha_- - \beta_- k^2) \hat{u}_k = g_-$$

$$\sum_{k=0}^N (-1)^k (\alpha_+ + \beta_+ k^2) \hat{u}_k = g_+$$

- ▶ Expressing  $\hat{u}_k^{(2)}$  and  $\hat{u}_k^{(1)}$  in terms of  $\hat{u}_k$  via the Chebyshev spectral differentiation matrices we obtain the following system

$$\mathbb{A} \hat{U} = \hat{F}$$

where  $\hat{U} = [\hat{u}_0, \dots, \hat{u}_N]^T$ ,  $F = [\hat{f}_0, \dots, \hat{f}_{N-2}, g_-, g_+]$  and the matrix  $\mathbb{A}$  is obtained by adding the two rows representing the boundary conditions (see above) to the matrix  $\mathbb{A}_1 = -\nu \hat{D}^2 + a \hat{D} + bI$ .

- ▶ When the domain boundary is not just a point (e.g., in 2D / 3D), formulation of the Tau method becomes somewhat more involved

- ▶ Consider the residual

$$R_N(x) = -\nu u_N'' + au_N' + bu_N - f,$$

where  $u_N(x) = \sum_{k=0}^N \hat{u}_k T_k(x)$

- ▶ Cancel this residual at  $N - 1$  GAUSS-LOBATTO-CHEBYSHEV collocation points located in the interior of the domain

$$-\nu u_N''(x_j) + au_N'(x_j) + bu_N(x_j) = f(x_j), \quad j = 1, \dots, N - 1$$

- ▶ Enforce the two boundary conditions at endpoints

$$\alpha_- u_N(x_N) + \beta_- u_N'(x_N) = g_-$$

$$\alpha_+ u_N(x_0) + \beta_+ u_N'(x_0) = g_+$$

Note that this shows the utility of using the GAUSS-LOBATTO-CHEBYSHEV collocation points

- Consequently, the following system of  $N + 1$  equations is obtained

$$\sum_{k=0}^N (-\nu d_{jk}^{(2)} + a d_{jk}^{(1)}) u_N(x_j) + b u_N(x_j) = f(x_j), \quad j = 1, \dots, N - 1$$

$$\alpha_- u_N(x_N) + \beta_- \sum_{k=0}^N d_{Nk}^{(1)} u_N(x_k) = g_-$$

$$\alpha_+ u_N(x_0) + \beta_+ \sum_{k=0}^N d_{0k}^{(1)} u_N(x_k) = g_+$$

which can be written as  $\mathbb{A}_c \mathbf{U} = \mathbf{F}$ , where  $[\mathbb{A}_c]_{jk} = [\mathbb{A}_{c0}]_{jk}$ ,  $j, k = 1, \dots, N - 1$  with  $\mathbb{A}_{c0}$  given by

$$\mathbb{A}_{c0} = (-\nu \mathbb{D}^2 + a \mathbb{D} + b \mathbb{I}) \mathbf{U}$$

and the **BOUNDARY CONDITIONS** above added as the rows **0** and  **$N$**  of  $\mathbb{A}_c$

- ▶ Note that the matrix corresponding to this system of equations may be **POORLY CONDITIONED** , so special care must be exercised when solving this system for large  $N$ .
- ▶ Similar approach can be used when the nodal values  $u(x_j)$ , rather than the Chebyshev coefficients  $\hat{u}_k$  are unknowns

- ▶ When the equation has **NONCONSTANT COEFFICIENTS**, similar difficulties as in the Fourier case are encountered (evaluation of **CONVOLUTION SUMS**)
- ▶ Consequently, the **COLLOCATION** (pseudo-spectral) approach is preferable along the guidelines laid out in the case of the Fourier spectral methods
- ▶ Assuming  $a = a(x)$  in the elliptic boundary value problem, we need to make the following modification to  $\mathbb{A}_c$ :

$$\mathbb{A}'_{c0} = (-\nu \mathbb{D}^2 + \mathbb{D}' + b\mathbb{I})U,$$

where  $\mathbb{D}' = [a(x_j)d_{jk}^{(1)}]$ ,  $j, k = 1, \dots, N$

- ▶ For the Burgers equation  $\partial_t u + \frac{1}{2}\partial_x u^2 - \nu\partial_x^2 u$  we obtain at every time step  $n$

$$(\mathbb{I} - \Delta t \nu \mathbb{D}^{(2)})U^{n+1} = U^n - \frac{1}{2}\Delta t \mathbb{D} W^n,$$

where  $[W^n]_j = [U^n]_j[U^n]_j$ ; Note that an algebraic system has to be solved at each time step

# Epilogue — Domain Decomposition

## ► Motivation:

- treatment of problem in **IRREGULAR DOMAINS**
- **STIFF PROBLEMS**

## ► **PHILOSOPHY** — partition the original domain $\Omega$ into a number of **SUBDOMAINS** $\{\Omega_m\}_{m=1}^M$ and solve the problem separately on each those while respecting consistency conditions on the interfaces

## ► **Spectral Element Method**

- consider a collection of problems posed on each subdomain  $\Omega_m$

$$\mathcal{L}u_m = f$$

$$u_{m-1}(a_m) = u_m(a_m), \quad u_m(a_{m+1}) = u_{m+1}(a_{m+1})$$

- Transform each subdomain  $\Omega_m$  to  $I = [-1, 1]$
- use a separate set of  $N_m$  **ORTHOGONAL POLYNOMIALS** to approximate the solution on every subinterval
- boundary conditions on interfaces provide coupling between problems on subdomains